

Synonym

CD86,B7-2,B70,CD28LG2,LAB72,MGC34413

Source

Human B7-2, His Tag(CD6-H5223) is expressed from human 293 cells (HEK293). It contains AA Leu 26 - Pro 247 (Accession # [AAH40261](#)).

Predicted N-terminus: Leu 26

Molecular Characterization

B7-2(Leu 26 - Pro 247)
AAH40261

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 26.2 kDa. The protein migrates as 40-63 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human B7-2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS




The purity of Human B7-2, His Tag (Cat. No. CD6-H5223) is more than 90% and the molecular weight of this protein is around 40-50 kDa verified by SEC-MALS.

[Report](#)


Discounts, Gifts,
and more!






Immobilized Biotinylated Human CTLA-4, Fc,Avitag (Cat. No. CT4-H82F3) at 5 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Human B7-2, His Tag (Cat. No. CD6-H5223) with a linear range of 0.01-2.5 µg/mL (QC tested).

Bioactivity-BLI



Loaded Human / Cynomolgus / Rhesus macaque CD28, Fc Tag (Cat. No. CD8-H525a) on Protein A Biosensor, can bind Human B7-2, His Tag (Cat. No. CD6-H5223) with an affinity constant of 6.8 µM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Human CTLA-4, Fc Tag (Cat. No. CT4-H5255) on Protein A Biosensor, can bind Human B7-2, His Tag (Cat. No. CD6-H5223) with an affinity constant of 1.9 µM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

Cluster of Differentiation 86 (CD86) is also known as B-lymphocyte activation antigen B7-2, is a type I membrane protein that is a member of the immunoglobulin superfamily, and is constitutively expressed on interdigitating dendritic cells, Langerhans cells, peripheral blood dendritic cells, memory B cells, and germinal center B cells. Additionally, B72 is expressed at low levels on monocytes and can be upregulated through interferon γ. CD86 is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD86 works in tandem with CD80 to prime T cells. Recent study has revealed that B7-2 promotes the generation of a mature APC repertoire and promotes APC function and survival. Furthermore, the B7 proteins are also involved in innate immune responses by activating NF-κB-signaling pathway in macrophages. CD86 thus is regarded as a promising candidate for immune therapy. CD86+ macrophages in Hodgkin lymphoma patients are an independent marker for potential nonresponse to firstline-therapy.

Clinical and Translational Updates



Human B7-2 / CD86 Protein, His Tag (MALS verified)

Catalog # CD6-H5223

