

Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

Human CD32a (131R) (Luc) Jurkat Reporter Cell


Catalog No.	Size
SCJUR-STF070	2 × (1 vial contains ~5×10 ⁶ cells)

• *Description*

The Human CD32a (131R) (Luc) Jurkat Reporter Cell was engineered to not only express the NFAT response element driving luciferase expressing systems, but also express the receptor full length human CD32a mutated to an Arginine (R) at amino acid 131 exhibiting a lower affinity for IgG2 isotypes compared to CD32a-131H, which can use to evaluate ADCP activity of antibodies in the presence of corresponding target cells. When co-cultured with a target cell and relevant antibody, the antibody simultaneously binds the target cell antigen and CD32a (131R) receptor on the surface of Human CD32a (131R) (Luc) Jurkat Reporter Cell, resulting in receptor clustering, intracellular signaling and NFAT-mediated luminescence.

Application

- Determination of ADCP activity induced by antibodies.



Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

• Cell Line Profile

Cell line	Human CD32a (131R) (Luc) Jurkat Reporter Cell
Host Cell	Jurkat
Property	Suspension
Complete Growth Medium	RPMI-1640 + 10% FBS
Selection Marker	Puromycin (5 µg/mL) + Hygromycin (20 µg/mL)
Incubation	37°C with 5% CO ₂
Doubling Time	16-20 hours
Transduction Technique	Lentivirus

• Materials Required for Cell Culture

- RPMI Medium 1640 (Gibco, Cat.No.11875-093)
- Fetal bovine serum (CellMax, Cat.No.SA211.02)
- Puromycin (InvivoGen, Cat.No.ant-pr-5b)
- Hygromycin B (Invitrogen, Cat.No.10687010)
- Complete Growth Medium: RPMI-1640 + 10% FBS
- Culture Medium: RPMI-1640 + 10% FBS, Hygromycin (20 µg/mL), Puromycin (5 µg/mL)
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, 430641)
- Cryogenic storage vials (SARSTEDT, 72.379.007)
- Thermostat water bath
- Centrifuge
- Luna cell counter (Logos Biosystems, LUNA- II)
- CO₂ Incubator (Thermo, 3111)
- Biological Safety Cabinet (Thermo, 1389)

Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

• *Recovery*

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the cap out of the water. Thawing should be rapid (approximately 5 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by spraying with 70% ethanol. All the operations from this point on should be carried out under strict aseptic conditions.
3. Transfer the vial contents to a centrifuge tube containing 4.0 mL complete growth medium.
4. Count viable cells and spin at approximately 1000 rpm for 5 minutes.
5. Discard the supernatant and resuspend the cell pellet in an appropriate amount of fresh complete growth medium. Adjust the cell density of the suspension to 1×10^6 viable cells/mL and transfer cells to an appropriate size vessel.
6. Incubate at 37°C with 5% CO₂ incubator

• *Subculture*

Adjust the cell density at 2×10^5 - 5×10^5 viable cells/mL by the addition of fresh culture medium or replacement of culture medium. Do not allow the cell density to exceed 3×10^6 cells/mL. T-75 flasks are recommended for subculturing.

- **Medium Renewal:** Add fresh culture medium every 3 to 4 days (depending on cell density)

• *Cryopreservation*

1. Count viable cells and harvest the cell suspension.
2. Centrifuge at 1000 rpm for 5 min at RT and resuspend cells in freezing medium to a concentration of 5×10^6 to 1×10^7 cells/mL.
3. Aliquot into cryogenic storage vials. Place vials in a programmable cooler or an insulated box placed in a -80°C freezer overnight, then transferring to liquid nitrogen storage.

• *Storage*

- **Product format:** Frozen
- **Storage conditions:** Liquid nitrogen immediately upon receipt

Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

• *Receptor Assay*




Fig1. Expression analysis of human CD32a (131R) on Human CD32a (131R) (Luc) Jurkat Reporter Cell by FACS. Cell surface staining was performed on Human CD32a (131R) (Luc) Jurkat Reporter Cell or negative control cell using PE-labeled anti-human CD32a antibody.

• *Application*




Fig2. ADCP response to anti-human CD20 antibody (RLU). Anti-human CD20 antibody-induced ADCP activity was evaluated using Human CD32a (131R) (Luc) Jurkat Reporter Cell in the presence of Raji cells that express CD20 endogenously. The EC₅₀ of anti-human CD20 antibody was approximately 0.037 µg/mL.

Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

Determination of ADCP activity




Fig3. ADCP response to anti-human CD20 antibody (Fold). Anti-human CD20 antibody-induced ADCP activity was evaluated using Human CD32a (131R) (Luc) Jurkat Reporter Cell in the presence of Raji cells that express CD20 endogenously. The max induction fold was approximately 88.

• *License Disclosure*

This reporter cell is provided for research use only. This license does not permit you to share, distribute, sell, sublicense, or otherwise make this reporter cell available for use to other laboratories, departments, research institutions, hospitals, universities, or biotech companies. The license does not permit modification of this reporter cell in any way. Inappropriate use or distribution of this reporter cell will result in revocation of the license. Modifications of this cell line, transfer to another facility, or commercial use of the cells may require a separate license and additional fees. AcroBiosystems does not warrant the suitability of this reporter cell for any particular use, and does not accept any liability in connection with the handling or use of this reporter cell.

Human CD32a (131R) (Luc) Jurkat Reporter Cell Development Service Data Sheet

• *Related Products*

Products

Human CD16a (158V) (Luc) Jurkat Reporter Cell Development Service

Human CD16a (158F) (Luc) Jurkat Reporter Cell Development Service

Human CD32a (131H) (Luc) Jurkat Reporter Cell Development Service

Human CD32b (Luc) Jurkat Reporter Cell Development Service

Human CD64 (Luc) Jurkat Reporter Cell Development Service

Cat.No.

SCJUR-STF067

SCJUR-STF068

SCJUR-STF069

SCJUR-STF071

SCJUR-STF072